

Contents lists available at ScienceDirect

Forest Ecology and Management



journal homepage: www.elsevier.com/locate/foreco

Litter share and clay content determine soil restoration effects of rich litter tree species in forests on acidified sandy soils



Ellen Desie^{a,*}, Karen Vancampenhout^b, Leon van den Berg^{c,d}, Bart Nyssen^{a,c}, Maaike Weijters^e, Jan den Ouden^f, Bart Muys^{a,*}

^a Division Forest, Nature and Landscape, KU Leuven, Celestijnenlaan 200E, Box 2411, B-3001 Leuven, Belgium

^b Department of Earth and Environmental Sciences, KU Leuven Campus Geel, Kleinhoefstraat 4, B-2240, Geel, Belgium

^c Bosgroep Zuid-Nederland, Huisvenseweg 14, 5591 VD Heeze, the Netherlands

^d Aquatic Ecology & Environmental Biology, Radboud University Nijmegen, Heyendaalseweg 135, 6525 AJ Nijmegen, the Netherlands

^e B-WARE Research Centre/Radboud University Nijmegen, Toernooiveld 1, 6525 ED Nijmegen, the Netherlands

^f Forest Ecology and Management Group, Wageningen University, P.O. Box 47, Wageningen, the Netherlands

ARTICLE INFO

Keywords: Litter quality Black cherry Soil restoration Clay content Soil acidification Nutrient cycling

ABSTRACT

Many West-European forests are located on degraded and acidified soils. Soil acidification has resulted in hampered ecosystem functioning and lower delivery of ecosystem services. Forest management, particularly the choice of tree species, can accelerate or counteract soil acidification by the quality of litter input. The positive impact of so called 'rich litter' on the soil nutrient status and belowground ecosystem functioning has already been evidenced in common gardens. Here, we evaluate the effect of the rich litter species black cherry (Prunus serotina Ehrh.) in mixed forest stands dominated by pedunculate oak (Quercus robur L.). We study the effects using a replicated set-up of 10 established forest stands (age 30 to 90) in Belgium, the Netherlands and Germany along an edaphic gradient in sandy soils on Pleistocene aeolian deposits. We hypothesize that black cherry has a positive effect on the soil nutrient status and aim to answer the following research questions: (i) does admixture of black cherry increase soil pH and base saturation? (ii) what proportion of rich litter admixture is needed in a poor litter matrix to observe significant improvement of the soil nutrient status? and (iii) does the magnitude of the rich litter effect interact with initial soil properties? The results of this study indicate that admixture of black cherry enhances the forest floor turnover and enriches topsoil chemical conditions significantly. Thickness of the litter layer decreases from a mean of 7 cm under oak to a mean of 4.5 cm under cherry and correspondingly base saturation increases to a maximum of 25%, NO₃⁻ concentration to 26 mg/mg and organic matter content to 8%. However, large shares of rich litter admixture (> 30% basal area) are needed to improve topsoil conditions. Moreover, we find that rich litter effects are more pronounced on sandy soils with higher fine particle (loam + clay) content. This suggests that the actual impact of restoration efforts in acidified forest soils is a product of the trinity "litter quality - litter share - site quality".

1. Introduction

Steady nutrient cycling between different compartments of ecosystems is a precondition for the provision of multiple ecosystem services (Lavelle et al., 2003). In that regard, soil acidification jeopardizes the long term functioning of forest ecosystems by altering the availability of critical macro- and micronutrients in the soil (Likens et al., 1996; Schaberg et al., 2001). Although this process occurs naturally in many forest soils, it has increased due to centuries of unsustainable land use, and further accelerated over the last century due to atmospheric acidifying deposition (Galloway, 2001). Soil acidification leads to the loss of the base cations calcium (Ca), magnesium (Mg) and potassium (K), and may increase concentrations of available aluminum (Al) and iron (Fe) to toxic levels (Ulrich and Sumner, 1991; Bowman et al., 2008), which negatively affects the vitality or growth of many plant and soil fauna. Despite coordinated international efforts to reduce atmospheric deposition of sulphur (S) and nitrogen (N) since the 1980s, current nitrogen deposition levels still exceed the critical load, i.e. the level below which no harmful effects can be expected (de Vries et al., 2014; Waldner et al., 2015). Especially in sandy soils, which are more vulnerable to degradation, the ensuing nutrient imbalance (i.e. base cation deprivation along with an overload of N) disturbs ecosystem

* Corresponding authors. *E-mail addresses*: ellen.desie@kuleuven.be (E. Desie), bart.muys@kuleuven.be (B. Muys).

https://doi.org/10.1016/j.foreco.2020.118377

Received 3 May 2020; Received in revised form 26 June 2020; Accepted 27 June 2020 0378-1127/@2020 Published by Elsevier B.V.

functioning and reduces the overall vitality of the ecosystem (Schaberg et al., 2001).

As such, forest vitality is far from recovered from the continuous input of acidity over the last century (Schmitz et al., 2019). For example, European oak forests are increasingly affected by acute oak decline, which has been linked to the mentioned changes in soil chemistry and nutrient status (Demchik and Sharpe, 2000; Brown et al., 2018). Oak remains one of the main timber producing species for the West-European wood industry, and simultaneously fulfills an important ecological niche with much associated biodiversity (Peterken, 1996). Increased mortality is not a new phenomenon, but arises when trees are under physiological stress and become more vulnerable to pathogens and tissue damage (Jung et al., 2000; Denman et al., 2010). In order to boost the resilience of West European oak forests, the current cation imbalances in the soil need alleviation and restoration efforts focusing on nutrient status.

To curb soil acidification and its consequences, many restoration strategies have been explored (Kreutzer, 1995; Hüttl and Schneider, 1998; Dumitru et al., 1999; Musil and Pavlicek, 2002). Trials with liming and application of rock dust show promising results, yet only intervene in the abiotic compartment of the forest ecosystem and do not couple belowground nutrient cycles with aboveground biomass, which is important on soils with low cation exchange capacity (CEC) (Formánek and Vranová, 2002). A long-term strategy, aimed at actively incorporating the aboveground compartment into nutrient cycling, is to admix tree species with a favorable litter composition to speed up nutrient cycling, promote more diverse soil communities and improve the soil chemical status (Finzi et al., 1998; Hommel et al., 2007a). In acidic edaphic conditions, litter rich in base cations (further: rich litter) promotes earthworm abundance which leads to more incorporation of organic matter in the soil (Muys et al., 1992). In turn, this leads to an improved nutrient binding capacity (higher CEC) and water holding capacity of the topsoil, creating higher resilience against future disturbance (e.g. acid input or drought). Previous studies evaluating tree species litter effects have identified rich-litter species (e.g. Tilia, Acer, Fraxinus and Prunus species) and evidenced their soil-enriching capacity on sandy soils (Desie et al., 2020; Reich et al., 2005; Mueller et al., 2015). However, these studies have evaluated pure tree species effects in common gardens or monoculture stands whereas species can behave differently in mixtures (Hättenschwiler et al., 2005). Tree species currently dominating in areas on sandy soils such as pine, oak, beech, larch and douglas fir all produce litter that is low in base cations (i.e. poorlitter species), therefore a more direct assessment of mixtures with such poor litter is necessary. Moreover, tree species effects differ relative to soil type (Verstraeten et al., 2018; Desie et al., 2019), which may explain why the few studies looking at rich-litter species admixture in poor-litter matrices on the soil nutrient status have reported mixed positive (Carnol and Bazgir, 2013; Aerts et al., 2017), none (Van Nevel et al., 2014) or even negative (Aerts et al., 2017) results. In terms of nutrient cycling, it is therefore important to evaluate whether the rich litter effect will still prevail when rich-litter trees are admixed in stands dominated by poor-litter tree species on a particular soil type, and how the share of that rich litter influences outcomes.

This study aims to evaluate whether admixture of a rich-litter species in pedunculate oak (*Quercus robur* L.) stands enhances the soil nutrient status of poorly buffered acidified sandy soils. We chose to study the admixture of black cherry (*Prunus serotina* Ehrh.) as we know from national forest inventories and our own observations that it is the only rich-litter species occurring in sufficient frequency (and as mature and dominant overstory tree) in admixtures of oak stands on relatively poor sandy soils in our study region. We wanted to make abstraction of the invasive species debate with regard to black cherry and use it as a model species for the evaluation of the impact of rich litter on nutrient cycles in forests on acidified sandy soils. We hypothesize that admixture black cherry has a positive effect on the soil nutrient status and is functional as a management measure for soil restoration. Particularly, we address the following research questions: (i) Does admixture of black cherry increase soil base saturation, pH and available N? (ii) How much admixture of rich litter is needed in a poor litter matrix to observe significant improvement of the soil nutrient status? And (iii) Does the magnitude of the rich litter effect depend on initial soil properties? These questions were answered using a replicated set-up established in 10 established (age 30 to 90), mixed forest sites in Belgium, the Netherlands and Germany along an edaphic gradient of sandy soils (texture ranging from 56% to 95% sand).

2. Materials and methods

2.1. Study region and sampling design

This study focused on mixed forest stands located in Northern Belgium, Southern Netherlands and the adjacent area in Germany (center 51° 17′ N, 5° 31′ E, altitude 30-80masl). The region is characterized by Pleistocene sandy aeolian deposits of variable thickness, locally admixed with sediments from marine or riverine origin (Kasse et al., 2007). Hence, soil textures vary from almost pure sand, over loamy or clayey sands to sandy loams (Van Ranst and Sys, 2000). The fraction of particles < 50 μ m can vary from ca. 50% to almost zero. The climate is temperate with a mean annual precipitation (MAP) of circa 800 mm and a mean annual temperature (MAT) of 10.5 °C (data provided by the Royal Meteorological Institute of Belgium). Ten sites were selected, based on the presence of a mixture of pedunculate oak (*Quercus robur* L.) and black cherry (*Prunus serotine* Ehrh.) in the upper canopy (Table 1) and along a gradient in soil texture.

In each mixed forest site, 4 replicates of 3 types of trees were selected: a dominant black cherry tree, a neighbouring oak tree under influence of black cherry (at a maximum of one tree height distance from the black cherry) and a reference oak tree without direct influence of dominant black cherry trees (Fig. 1). By selecting the reference trees in the same forest stand on the same site (maximum 230 m distance from oak under influence) and by evaluating historical maps, we assured that all selected trees grew under the same environmental conditions (climate and topography) and had the same land-use legacy. By sampling both cherry and oak in the same forest stand confounding factors were limited as to assure that the actual differences reported can be appointed to tree species effects and not initial differences in site conditions. We may even underestimate black cherry influence as, because of the design, some reference oak trees were under minor influence of black cherry. In four sites (Genk, Mol, Walbeck, t'Zand) we could not find four dominant black cherry trees present that met our selection criteria, therefore we sampled two oak trees under influence of the same dominant cherry present. In total we selected and sampled 115 trees.

2.2. Sampling and laboratory analysis

2.2.1. Sampling

Under each selected tree, soil samples were taken and humus descriptions were made in July 2017. We sampled around each selected tree in all wind directions at a distance of 1/3 crown radius from the stem. For each selected tree the humus layer was described three times following the European humus reference base (Zanella et al. 2014). We measured the thickness of the OL, OF and OH layer separately and determined the humus type, here expressed as the humus index (HI) (Ponge et al., 2002). Further analysis of these data was based on the mean of the 3 replications (median for humus index). Around each selected tree, five bulk mineral soil samples were taken from two depths (0–10 cm and 20–30 cm mineral soil depth for topsoil and subsoil samples respectively, where 0 cm depth starts directly under the forest floor layer at the top of the mineral soil), and merged in one composite sample per depth for chemical analysis. The forest floor layer itself was not sampled. At each site an augering was performed and the soil type

 Table 1

 Characteristics per site located in Belgium (B), the Netherlands (NL) or Germany (D). Texture and soil type were determined per site (group-level variables). The mean of subsoil CEC, OM, N, P, BA and species composition was calculated based on all selected trees per site (population-level variables). The earliest indicated of forest on available maps is indicated as 'forested since' (note this is not the stand age). Sites are ranked based on

E. Desie, et al.

IIICLEASHIG SAHU COHEIL											
Site	Texture			Soil type	Topsoil (0–10 d	(m					Subsoil (20–30 cm)
	clay	Silt	Sand	reference group	Hq	BS	Al	$NO_{3}^{-} + NH_{4}^{+}$	CEC	MO	$NO_3^- + NH_4^+$
	(%)	(%)	(%)			(%)	mg/kg	mg/kg	(meq/100 g)	(%)	(mg/kg)
Genk (B)	12.7	30.5	56.8	Arenosol	3.11 ± 0.10	33.77 ± 13.95	19.53 ± 6.86	41.56 ± 21.67	4.07 ± 1.35	7.09 ± 2.92	5.26 ± 2.48
Veldhoven (NL)	3.6	29.0	67.4	Anthrosol	3.20 ± 0.11	23.86 ± 13.50	42.80 ± 6.60	30.72 ± 22.46	3.60 ± 0.88	5.58 ± 1.84	4.43 ± 2.20
Walbeck (D)	6.7	24.5	68.8	Arenosol	2.86 ± 0.06	22.75 ± 9.30	70.42 ± 19.37	67.16 ± 26.44	7.39 ± 2.23	16.44 ± 4.79	8.35 ± 3.92
As (B)	5.5	23.6	70.9	Anthrosol	3.23 ± 0.09	35.20 ± 6.79	18.57 ± 4.61	36.40 ± 14.51	3.53 ± 0.63	6.26 ± 0.93	5.00 ± 2.54
Loon op zand (NL)	2.0	19.8	80.4	Podzol	3.05 ± 0.12	12.56 ± 6.11	38.18 ± 9.10	12.78 ± 5.02	4.08 ± 1.28	5.82 ± 2.10	4.50 ± 4.32
Someren (NL)	2.0	13.2	86.6	Arenosol	3.22 ± 0.14	13.56 ± 3.86	36.84 ± 5.78	17.28 ± 5.48	3.56 ± 0.78	4.55 ± 1.01	6.07 ± 4.17
Grashoek (NL)	1.7	9.1	89.2	Podzol	2.99 ± 0.06	16.51 ± 11.95	33.60 ± 12.36	17.43 ± 6.14	3.82 ± 0.79	5.01 ± 1.68	7.00 ± 3.69
Mol (B)	1.1	5.3	93.9	Arenosol	3.33 ± 0.30	32.19 ± 12.52	28.71 ± 10.74	46.50 ± 17.10	5.87 ± 1.17	9.97 ± 1.26	9.05 ± 4.34
t'Zand (NL)	1.3	3.9	94.8	Arenosol	3.15 ± 0.10	21.50 ± 10.80	30.43 ± 4.23	7.42 ± 3.67	2.55 ± 0.65	3.02 ± 1.72	2.82 ± 1.89
Hoogstraten (B)	1.0	4.5	95.1	Podzol	2.83 ± 0.14	8.70 ± 3.14	39.60 ± 13.00	16.01 ± 7.29	4.77 ± 0.78	5.93 ± 1.20	9.02 ± 12.69
Site	Subsoil (20–30	cm)			Species composi	tion (%)					Forested
	CEC	MO	Total P	Basal area	Quercus robur	Prunus Pinu serotina	us sylvestris Betula	pendula Fagus sylvatica	Quercus rubra	other	sınce (according to historical
	(meq/100 g)	(%)	(mg/kg)		(%)	(%) (%)	(%)	(%)	(%)	(%)	maps)
Genk (B)	1.65 ± 0.40	2.06 ± 0.68	263.24 ± 39.22	2.42 ± 0.40	38	12 46	1				1971
Veldhoven (NL)	1.47 ± 0.32	1.91 ± 0.25	262.9 ± 30.15	1.71 ± 0.77	46	19 3	6			6	1850
Walbeck (D)	2.87 ± 0.64	3.98 ± 1.28	184.01 ± 0.54	2.05 ± 0.33	60	24 6	1				1850
As (B)	1.61 ± 0.12	2.5 ± 0.37	183.23 ± 23.90	2.23 ± 0.61	52	28 10	8				1971
Loon op zand (NL)	1.96 ± 0.48	2.08 ± 0.92	39.54 ± 7.32	$1,82 \pm 0.35$	54	13	8	8			1988
Someren (NL)	2.19 ± 1.17	2.23 ± 0.79	184.99 ± 71.75	1.71 ± 0.24	56	23 3	13			S	1983
Grashoek (NL)	3.85 ± 1.25	4.11 ± 1.37	71.75 ± 37.49	1.95 ± 0.47	27	19 19	32				1926
Mol (B)	2.64 ± 1.23	3.48 ± 0.82	135.36 ± 61.63	2.03 ± 0.39	57	15 15	8		2		1971
t'Zand (NL)	1.36 ± 0.144	2.96 ± 4.64	28.77 ± 7.83	2.39 ± 0.42	50	15 27	L	·	,		1899
Hoogstraten (B)	3.49 ± 1.08	3.79 ± 1.00	09.04 ± 17.90	1.09 ± 0.29	51	34	n	Т	T		1840

3

was described according to the FAO guidelines for soil description (FAO, 2006). Soil classifications were executed on each site according to WRB guidelines (IUSS Working Group WRB, 2015). An additional soil sample of the subsoil (C-horizon) was sampled on each site for texture analysis.

In autumn 2018 we placed one litter trap (1 m high, 0.2 m² circular surface area) under each selected tree at a distance of 1/3 crown radius from the stem. Per site we placed all traps at the same orientation (dominant wind direction) from the tree. Litter traps were emptied 3 times between October and December. The collected material was oven dried at 60 °C and weighed per litter category (leaf litter of oak, cherry, conifers and other broadleaved, and non-foliar litter) and per collection date (Fig. S2). Data for selected trees with damaged litter traps (n = 26) were omitted.

In the winter of 2018–2019 we mapped the forest structure of all sites using the FieldMap instrument (FieldMap, IFER, Czech Republic). All trees with a diameter at breast height (DBH) higher than 15 cm (Vannoppen et al., 2020) and within a radius of 15 m around the selected tree were spatially mapped and species, DBH and height of the tree were included in the map.

2.2.2. Laboratory analysis

Soil pH, NO3⁻ and NH4⁺ concentration were determined in salt extracts (after mixing fresh soil (17.5 g) with 50 ml 0.2 M NaCl solution). The $pH_{(NaCl)}$ was measured immediately after extraction using a combined pH electrode (radiometer and a TIM840 pH meter). NO₃and NH4⁺ concentrations were determined colorimetrically with a Seal auto-analyser III, using salicylate, hydrazin sulphate and ammoniummolybdate/ascorbic acid reagent, respectively (Grasshoff and Johannsen, 1977; Technicon, 1969). Acid extractable element concentrations (Al, Ca, Fe, K, Mg, Mn, S, Si, P, Zn) of soil and litter samples were determined by digesting 200 mg of dried (24 h, 70 °C) and homogenized (by mortar) sample in 4 ml concentrated HNO3 and 1 ml 30% H2O2 (Milestone microwave MLS 1200 Mega) (Kingston and Haswell, 1997). Cation Exchange Capacity (CEC) and base saturation were determined by mixing an amount of dry soil equivalent of 5 g fresh soil in 200 ml 0.2 M SrCl₂ (Liu et al., 2001). All the soil extracts were measured with ICP, as mentioned above. Base saturation was calculated as the sum of exchangeable Ca^{2+} , Mg^{2+} and K^+ (in terms of charge equivalents) divided by the CEC and expressed as %. Soil and leaf litter total nitrogen (N) and carbon (C) concentrations were measured with a CNS analyzer (Model NA 1500; Carlo Erba Instruments, Milan, Italy). Soil organic matter content was determined by weighed loss-on-ignition after burning samples at 550 °C for a minimum of 6 h (Schulte and Hopkins, 1996). Soil texture was analyzed by laser diffractometry using a laser diffraction particle size analyzer - LS 13 320) (Buurman et al., 2001). The fine particle content is based on the sum of the clay content and loam content.

2.3. Data analysis

2.3.1. Stand characteristics

Based on the collected stand structure data, following variables were calculated for each selected tree. The basal area (BA) of all neighbors was calculated as follows: $\sum_{i=1}^{n} 0.25\pi (d_i)^2$ in a 15 m radius with d_i the DBH of the ith tree. Competition was calculated using $\sum_{i=1}^{n} di/(d \times dist_i)$ according to Contreras et al. (2011) with dist_i representing the distance between the selected tree with diameter d and the competing tree i with diameter d_i. The influence of black cherry (further called Prunus influence) on each selected tree was calculated using the BA of black cherry in a 15 m radius around the selected tree (including the selected tree itself) where the BA area was weighted

based on the relative distance to the selected tree $(dist_j)$ and relative to the total amount of neighbors (of any species) in a 15 m radius:

Prunus influence =
$$\frac{\sum_{i}^{l} \left(\left(1 - \frac{dist_{i}}{15m} \right) BA \quad cherry_{i} \right)}{\sum_{i}^{l} \left(\left(1 - \frac{dist_{i}}{15m} \right) BA \quad neighbor_{j} \right)}$$

2.3.2. Statistical analysis

Descriptive statistics were executed for most tree-species independent soil properties per site (Table 1). The effect of Prunus influence on different soil properties was tested by means of mixed models with site as a random effect using the package nlme in R. The normality of the residuals and their relation to the fitted values were evaluated graphically. When Prunus influence was a significant predictor in the mixed model, it is portrayed by a solid line in the Figs. 2, 3, S3 and S7, other non-significant relations are portrayed by dotted lines. Additionally, clay content (or fine particles content respectively) was included in the mixed models twice: as a main effect and in interaction with Prunus influence. Because clay content is a group-level variable, i.e. only measured per site, and the response soil properties are population-level variables, i.e. measured under each selected tree (12 times per site), we are dealing with multilevel data. This implies that caution is necessary when interpreting the P-values of the interaction effect with clay content (Qian et al., 2010). Significance of the predictors was tested by a type II analysis of variance (anova) using the package car in R. All predictors were standardized (to a of mean of 0 and sd of 1) so that the standardized coefficients (Table 2) can be interpreted easily. All analyses were performed in R version 3.4.4 (R Core Team, 2019).

3. Results

3.1. Research design

Texture of the study sites ranges from sandy loam to sand (56% -95% sand) (Table 1). Soils belong to three different WRB Soil Reference Groups, i.e. Anthrosols, Arenosols and Podsols, which is representative for the typical gradient in soil properties in sandy sites in Belgium and the Netherlands. In terms of land-use history, some sites were afforested 170 years ago whereas others only recently (30 years ago after a period of agricultural use or heathland cover). Subsoil P and N show considerable variation among sites ranging from 28.77 to 263.24 mg/kg DW for total P and 2.82 to 9.05 mg/kg DW for the sum of ammonium and nitrate. The overstory tree species composition varied over the 10 sites, with the respective proportion of pedunculate oak ranging from 27% to 60% and that of black cherry from 12 to 38% (based on basal area). Evaluation of the design showed that the weighted influence of black cherry, based on basal area share in a 15 m radius around the selected tree, increased from the reference tree (mean $2\% \pm 3\%$) to the oak under influence (mean $30\% \pm 14\%$) to the soil under the black cherry tree itself (mean $42\% \pm 16\%$) (Fig. S1).

3.2. Prunus influence

3.2.1. Litter layer and humus classification

Accumulation in the OF and OH layer decreased with increasing Prunus influence (P < 0.001) (Fig. 2a and Fig. S4). Moreover, we found significant differences between each type of selected tree: the thickness of the OF-OH layer was larger under the reference oak trees compared to the oaks under influence (Fig. S5). The humus type did not significantly change with increasing Prunus influence (P = 0.44) (Fig. 2b).

3.2.2. Topsoil chemistry

Topsoil base saturation increased significantly with increasing

E. Desie, et al.



Fig. 1. (a) Study design and (b) study region located over Belgium, the Netherlands and Germany. The sampling design exists of 3 types of target trees: a dominant black cherry tree, an oak under influence of black cherry and a reference oak without direct influence of mature black cherry trees. Per site four trees of each type are selected (N = 12 per site). The 10 mixed forest sites are indicated by purple dots. Photos of site in Walbeck (top right) and Mol (bottom right). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Prunus influence (P = 0.001) (Fig. 3a). This was mirrored in the differences between all types of selected trees: mean base saturation increased from 19 \pm 10% under the reference oak to 21 \pm 13% under the oak under influence and 25 \pm 15% under the black cherry tree. Only base saturation under the cherry tree differed significantly from under the reference oak (Fig. S6). An increase in basal area of black cherry in a 15 m radius from 0% to 80% black cherry (corrected for distance) corresponds with an increase in topsoil base saturation from 22% to 25% (Fig. 3 and Table 2). This trend was not significant for topsoil $pH_{(NaCl)}$ (P = 0.15) and topsoil aluminum (P = 0.09) (Fig. 3d). Topsoil NO₃⁻ increased significantly with increasing Prunus influence (P = 0.001) (Fig. 3b, Table 2). No trend was found for topsoil NH₄⁺ (P = 0.38) (Fig. 3e). Topsoil CEC showed no relation with Prunus influence (P = 0.14), however a positive linear relation was found with topsoil organic matter (OM) concentration (P = 0.006) (Fig. 3f and c). For subsoil chemistry we only found a marginally significant relation between NO_3^- concentration and Prunus influence (P = 0.05) (Fig. S7).

3.3. Mediation by clay content

Clay content has, additionally, a significant positive effect on topsoil base saturation, topsoil OM and topsoil NO_3^- . Finally, we found

significant positive interactions between Prunus influence and clay content for thickness of the OF-OH layer, topsoil base saturation, topsoil OM and topsoil NO_3^- indicating the greater impact of prunus on sites with a higher clay content. We found the same significant interaction effects with the total fine fraction (silt + clay content, reported in Table S3).

We evaluated the slopes of the topsoil – Prunus influence relation per site (Fig. S8) and plotted them as function of clay (Fig. 4 left) and fine particles content (clay + silt) (Fig. 4 right) to graphically assess the interaction effect, i.e. how texture magnifies the cherry effect (Fig. 4, Fig. S8).

To evaluate the correlation between texture and Prunus influence we investigated the potential link with base cation concentration of the cherry litter Fig. 5. Yet, base cation content of the leaf litter of all selected black cherries per site is not significantly related to clay content or fine particle content, even when accounting for the outlying site Mol (Fig. 5). When evaluating litter quality of the cherry trees we found significant relations between litter base cations and topsoil base saturation (P = 0.005) and topsoil pH (P < 0.001) (Fig. S9).

Table 2

Standardized coefficients, standard deviations and P-values of fixed effects (intercept, Prunus influence, clay and prunus*clay interaction) of mixed models accounting for site as a random effect explaining different topsoil response variables.

Response	Intercept	Prunus influence	Clay	Interaction
OF + OH layer Topsoil base saturation Topsoil pH _(NaCl) Topsoil Al Topsoil OM Topsoil CEC Topsoil NO ₃ ⁻ Topsoil NH ₄ ⁺	$\begin{array}{l} 5.47 \ (\ \pm \ 0.82)^{P < \ 0.001} \\ 22.28 \ (\ \pm \ 2.49)^{P < \ 0.001} \\ 3.10 \ (\ \pm \ 0.05)^{P < \ 0.001} \\ 35.62 \ (\ \pm \ 4.90)^{P < \ 0.001} \\ 6.99 \ (\ \pm \ 1.20)^{P < \ 0.001} \\ 4.32 \ (\ \pm \ 0.45)^{P < \ 0.001} \\ 22.83 \ (\ \pm \ 4.82)^{P < \ 0.001} \\ 6.41 \ (\ \pm \ 1.04)^{P < \ 0.001} \end{array}$	$\begin{array}{l} -1.09 (\pm 0.21)^{P < 0.001} \\ 3.11 (\pm 0.93)^{P = 0.001} \\ 0.02 (\pm 0.01)^{P = 0.15} \\ 1.70 (\pm 1.01)^{P = 0.09} \\ 0.58 (\pm 0.20)^{P = 0.006} \\ 0.16 (\pm 0.11)^{P = 0.14} \\ 4.22 (\pm 1.22)^{P < 0.001} \\ 0.28 (\pm 0.48)^{P = 0.66} \end{array}$	$\begin{array}{l} -0.97 (\pm 0.75)^{P=0.19} \\ 5.40 (\pm 2.38)^{P=0.02} \\ -0.009 (\pm 0.05)^{P=0.85} \\ 0.71 (\pm 4.26)^{P=0.86} \\ 1.89 (\pm 0.99)^{P=0.06} \\ -0.02 (\pm 0.40)^{P=0.94} \\ 15.99 (\pm 4.36)^{P<0.001} \\ -0.86 (\pm 1.01)^{P=0.39} \end{array}$	$\begin{array}{l} -0.59 (\pm 0.27) \stackrel{P=0.03}{} \\ 2.55 (\pm 1.22) \stackrel{P=0.03}{} \\ -0.01 (\pm 0.01) \stackrel{P=0.53}{} \\ -1.06 (\pm 1.33) \stackrel{P=0.42}{} \\ 0.68 (\pm 0.27) \stackrel{P=0.015}{} \\ 0.25 (\pm 0.14) \stackrel{P=0.08}{} \\ 4.32 (\pm 1.60) \stackrel{P=0.008}{} \\ 0.53 (\pm 0.63) \stackrel{P=0.39}{} \end{array}$



Fig. 2. (a) Thickness of the OF + OH layer and (b) humus type as a function of Prunus influence. The type of tree is indicated by the color and shape of the circles: reference oak (light green – fill), oak under influence (light green – no fill), black cherry (dark pink - fill). P-values for the variable 'Prunus influence' determined in a mixed model (Table 2) accounting for site effects are indicated in the top right corner. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

4. Discussion

4.1. Rich litter effect of Prunus serotina?

4.1.1. Litter quality

Previous studies illustrated the difference in forest floor thickness and humus form between different tree species in monoculture stands on the same site, including the large difference in litter quality between black cherry and pedunculate oak (von Wendorff, 1952; Vanderhoeven et al., 2005; Dassonville et al., 2008; Desie et al., 2020). Although the effect of tree species, for example litter quality, can be different in mixtures compared to pure stands (Hättenschwiler et al., 2005), we found that the effect of black cherry reported for monocultures, is maintained in mixed oak stands on acidic sandy soils. Increasing Prunus influence is significantly and negatively related to litter accumulation in the forest floor. This effect, also reported by Lorenz et al. (2004a), can be attributed to the promotion of microbial and faunal activity through the provision of base cation rich litter and subsequent improved decomposition (Hobbie et al., 2006; Reich et al., 2005). In this study we make the assumption that the positive effects are linked to the high litter quality of cherry, as we did not measure oak litter quality and cannot directly evidence the higher quality of litter of cherry compared to oak. This high litter quality was also the main reason for its massive introduction in West-Europe (Lorenz et al., 2004a).

Our study indicates that the effect of admixture of black cherry on topsoil chemistry is multiple. Admixture of cherry in acidified oak stands enhanced the nutrient status of the soils (both NO37 concentration as well as base saturation increase in the topsoil). The positive relation with topsoil base saturation can be explained by the higher base cation concentrations of fresh black cherry litter and the related accelerated turnover and incorporation of organic matter (Lavelle et al., 2004; Reich et al., 2005). Higher NO_3^- concentrations in proximity of black cherry are probably a consequence of the promoted nitrification explained by improved edaphic conditions for nitrifying microbes. Indeed an earlier study has found increased nitrification rates with increased soil pH and buffering (Ste-Marie and Paré, 1999). Despite the significant enrichment under black cherry, base saturation does not increase above the 30% threshold that is linked to a shift in soil buffering domain from a state dominated by Al to a state dominated by base cations (Vitousek and Chadwick, 2013; Desie et al., 2019). Soil pH changes when one buffering mechanism (for example Al buffering) is replaced by another (for example base cation buffering). In acid soils with pH below 4.5, the exchange complex is saturated with H⁺, Fe and Al ions, and soil pH will only respond markedly if base saturation can be raised above 30%. The soils in our study remain in the Al buffering range. As a result, the pH of the topsoil did not increase significantly



Fig. 3. Topsoil variables as a function of Prunus influence: (a) base saturation (%), (b) NO_3^- concentration (mg/kg DW), (c) organic matter content (%), (d) pH_(NaCl), (e) NH_4^+ concentration (mg/kg DW) and (f) CEC (meq/100 g). The type of tree is indicated by the color and shape of the circles: reference oak (light green – fill), oak under influence (light green – no fill), black cherry (dark pink - fill). Significant relations are represented by full lines whereas relations that are not significant are indicated by dotted lines. P-values for the variable 'Prunus influence' determined in a mixed model (Table 2) accounting for site effects are indicated in the top right corner. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)



Fig. 4. Magnitude of the rich litter effect (mean and se), i.e. the slope of the relation with Prunus influence, as a function of clay content (left) and fine particles content (clay + silt) (right). The magnitude is expressed as the slope of the topsoil base saturation or NO_3^- concentration – Prunus influence relation per site.

with black cherry admixture.

Although organic matter turnover and incorporation increased under black cherry, illustrated by the relation with topsoil OM, a significant relation with topsoil CEC was not found. We expected the CEC to increase as OM significantly contributes to the number of exchange places, especially in sandy soils with low CEC_{clay} (Gruba and Mulder, 2015). The lack of effect on CEC can potentially be explained by the pHdependent charge of the OM, i.e. OM will only contribute significantly to the CEC if the pH increases.

4.1.2. Litter share

Although we found significant effects of rich litter admixture, the effects are small in absolute terms. This corresponds with a previous study by Van Nevel et al. (2014) who found that dense understory shrub layers (with up to 90% cover) of rich litter species, among which black cherry, had no impact on topsoil chemistry of acidified sandy soils as the contribution to total litterfall was insufficient. They suggest that an improved overstory tree species selection would have more potential to improve topsoil conditions as compared to dense shrub layers of rich litter species. Indeed, our study corroborates these propositions, because we find significant positive effects of mature cherry trees on the soil conditions, which can be explained by the longer time period that rich litter effect has been active and the larger volume of litter produced by a mature tree. Moreover, the relative amount of rich-litter tree species needed in the overstory will depend on species-specific concentrations of base cations in the leaves. For example admixture of



Fig. 5. Base cation content of the black cherry litter (mean and se) per site as a function of clay content (left) and fine particles content (clay + silt) (right). Mol is (graphically) identified as an extreme value.

mature *Betula pubescens* trees (up to 63% of BA) proved insufficient to improve mineral soil conditions in *Picea abies* stands (Brandtberg et al., 2000). This can be explained by the relatively lower litter quality of birch, i.e. lower litter base cation concentrations, in comparison to other rich litter tree species (Desie et al., 2020), emphasizing that the community weighted mean of the litter composition remains the essence of soil nutrient status restoration.

We took both litter quality and the share of rich litter admixture into account by using mixed stands with mature, dominant black cherry trees, a tree species that has high concentrations of base cations in its litter (Desie et al., 2020). In our stands, an increase in basal area of black cherry from 0% to 80% black cherry (corrected for distance) translated in an actual increase of share in litterfall from 10% to 40% of black cherry, and corresponds with a small increase in topsoil base saturation from 22% to 25%. The 10% black cherry admixture in the litterfall of plots with 0% black cherry basal area shows that our basal area calculations cause an underestimation of black cherry litter share, because the basal area only included trees with a diameter higher than 15 cm, while the smaller diameter classes (< 15 cm) were almost all black cherry. Despite the fact that the actual leaf litter share in stands with high basal area remains limited, base saturation increased significantly. It is however clear that on poorly buffered sandy soils large amounts of admixture of black cherry (i.e. large concentrations of base cations) are needed to significantly impact the forest floor thickness and the soil base saturation and nutrient status. It remains to be seen if a longer period of rich litter influence will be able to increase base saturation further.

4.1.3. Limitations of the study

We found that cherry admixture has a positive effect on the forest floor and topsoil chemistry in mixed forest stands. Since we did not sample and measure litter quality of the oaks, we cannot directly evidence that the improved topsoil chemistry is a consequence of the richer litter of cherry. However, there is an extensive literature base that reports the high quality litter of black cherry (von Wendorff, 1952; Vanderhoeven et al., 2005; Desie et al., 2020) and explores the potential of this species for soil restoration (Carnol and Bazgir, 2013; Van Nevel et al., 2014). Therefore it is the most probable explanation. Furthermore, the base cation concentrations of cherry litter reported in this study may be an underestimation due to the long period between collection dates and potential leaching of mobile compounds from the litter in the traps. Secondly, we extended our litter trap mass data, which had multiple missing values, with more precise and quantitative basal area measurements of the overstory composition, which can be used as a proxy for litter contributions (Jonard et al., 2006; Nickmans et al., 2019). The correlation of both measures is illustrated in Fig. S3 and Table S2. We have made the assumption that the contribution of cherry in basal area corresponds with the contribution of litter to the forest floor. However, it should be addressed as a limitation of our study that we do not directly use litter share based on litter mass measurements.

4.2. Impact on the aboveground ecosystem compartment

Current forest management strategies are increasingly focused on boosting tree diversity in order to have higher insurance in a future shaped by increased frequency of disturbances (Paquette et al., 2018). Admixture of rich litter tree species offers an additional incentive through its ability to restore acidified, nutrient-imbalanced forest soils and is hypothesized to improve the overall vitality of the forest. The accelerating effect on nutrient cycling by rich litter species, such as black cherry, has been reported before (Lorenz et al., 2004; Vanderhoeven et al., 2005; Dassonville et al., 2008). Yet, whether this increased availability translates in higher uptake of nitrogen by the neighboring oak trees or, contrary, is short-circuited to the black cherry tree via its dense and superficial root system, remains an important question to be answered. The competitive behavior of black cherry could be appointed to the rooting system. Hence, subsequent research should evaluate the relative importance of litter and root dynamics of black cherry and whether this is also context dependent. In this study we merely focused on the litter-soil pathway. In terms of nutrient cycling, studies report diverse and even contradicting results: from increasing nutrient concentrations and pH (Lorenz et al., 2004; Vanderhoeven et al., 2005; Dassonville et al., 2008) to negatively affecting the belowground nutrient status (pH and N) (Starfinger et al., 2003) or having negative effects on neighboring native species and their foliage nutrient uptake (Aerts et al., 2017). Aerts et al. (2017) reported lower foliar nutrient concentrations of beech and oak in the close presence of black cherry, whereas higher foliar P concentration in pines were found in a different edaphic setting.

Finally, we want to point out that in this study on the management intervention "*admixing rich litter tree species*" black cherry served merely as a model species, that in the implementation of restoration can be replaced by other, native rich litter tree species better adapted to the management goals. In our previous study (Desie et al., 2020) we highlighted the potential of *Tilia, Acer* and *Alnus* as potential rich litter species for soil restoration. Moreover, further research should tackle the feedback of increased nutrient availability to the aboveground ecosystem and elucidate which species, in what conditions, have beneficial effects on the vitality of other trees.

4.3. Interaction with edaphic factors

We found significant relations between soil texture and topsoil base saturation and topsoil nitrate concentration. More intriguing, however, is the significant interaction between texture and rich litter effects: the higher the fraction of clay or fine particles (clay and silt combined) the greater the positive impact of admixing black cherry on topsoil conditions. This positive interaction cannot be explained by the higher litter quality of black cherry on sites with finer textures as we found no significant relation between litter base cation content and texture, implying that there is no feedback via base cations in the litter. We did find correlations between litter base cation content and topsoil base saturation and topsoil pH. Moreover, the high base cation content of litter from the site Mol can be explained by the input of base cations via Ca rich groundwater (neighboring a canal with Ca-rich water), which also explains the high subsoil base saturation in this site.

In our observations, clay content (or fine particle content) amplifies the positive effect of higher litter quality. Seemingly in contrast, Verstraeten et al. (2018) and Desie et al. (2019) concluded from their results that restoration may be more difficult in soils with a high clay content due to Al saturation of the exchange complex with aluminum and thus restoration on sandy soils is more feasible. These opposing trends in the relation between clay content and tree species litter effects can be explained by the context:

In soils with high CEC, Al saturation is typically difficult (or even impossible) to overcome since the total amount of aluminum sorbed on exchange sites is too high to be replaced by base cations through litter input. For agricultural settings a maximum value of CEC 24 meq/100 g clay is set (Driessen, 2001). Below this threshold restoration via litter input has more potential.

On the other hand, in soils with very low CEC values, such as our current study (subsoil CEC ranging between 1.36 and 3.85 meq/100 g soil DW with an average of 2.30 meq/100 g soil DW), the added benefit of a larger fine fraction in terms of soil fertility, SOM stabilization, aggregation, weatherable reserve and water holding capacity most likely is more determining than Al saturation on the exchange complex, explaining the positive relation with clay content.

Hence, the direction of the interaction with clay is context dependent, as biogeochemical equilibria in soils display considerable pedogenic inertia (so-called soil process domains; Ulrich and Sumner, 1991; Vitousek and Chadwick, 2013), interchanged by steep thresholds when one mechanism is exhausted and replaced by another (Chadwick and Chorover, 2001). Depending on the acid buffering capacity (which is arguably proxied by CEC) and the distance to a pedogenic threshold (proxied by base saturation), tree species can therefore have extensive, limited or no effect on soil acidification. Acidified soils that consist of almost pure sand or, contrary, acidified soils with a high clay content are opposite extremes (in CEC) that are trapped in so called pedogenic inertia and therefore very hard to affect by aboveground litter quality input.

Concluding, there is a window of opportunity for restoration in terms of site quality. In sandy soils (with low CEC), yet with considerable contribution of fine fractions (clay and silt), the admixture of rich litter trees can have maximum impact, i.e. around a pedogenic threshold in acid buffering small changes can have large impact. Hommel et al. (2007a) indicated a requirement of > 15% loam for admixture of rich litter to be successful. Van Nevel et al. (2014) also attribute the absence of rich litter effects to the high sand percentage of their study site. Nonetheless, under pure rich litter stands on poorly buffered sandy soils, we found significant positive effects (Desie et al 2020). This suggests that the ultimate result of restoration efforts is a product of the trinity litter quality – litter share – clay content. Hence, our study is emphasizing once more the importance of taking into account the boundary conditions determined by soil type when making forest management decisions.

5. Conclusion

Our study provides evidence that admixture of rich litter trees, in this case black cherry, in pedunculate oak stands leads to less accumulation of organic material in the forest floor and improves topsoil chemical conditions (base saturation, NO3⁻ and organic matter content) significantly. Hence, we accept our hypothesis that rich litter admixture can be used as a management measure for soil restoration. However, although the conditions improved, they did change not to such an extent that it does induce a regime shift to a more favorable soil process domain. We found that large shares of rich litter admixture (> 30% basal area) are needed to improve soil conditions and that such rich litter effects are more pronounced on sandy soils with higher clay content. Optimization of tree species selection as a management tool should focus on the regeneration of tree species with rich litter (i.e. elevated base cation concentrations), augmenting the relative share of such tree species in the forest overstory composition and targeting efforts to sites where pedogenic inertia does not limit the potential, i.e. soils with either very low or very high clay contents and corresponding CECs. Admixing rich litter holds promise and may ultimately restore nutrient cycling in forests of acidified sandy soils.

Author contributions

ED, KVC, LVDB, BN, JDO, BM designed the study; ED, MW collected the data; ED analyzed the data; All authors contributed to the interpretation of the results; ED compiled the manuscript; All authors contributed critically to the drafts and gave final approval for publication.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank *Bosgroep Zuid Nederland* for the coordination of the applied umbrella project and the Dutch province *Noord-Brabant* for funding. We would like to acknowledge the support of B-Ware research

centre regarding the laboratory analysis and Eric Van Beek, Remi Chevalier, Fien Decoster, Simon Baert, Bjorn Rombouts, Elisa Van Cleemput, Stef Boogers and Jan Vanstockem for their support during the data collection. Finally, we thank Koenraad Van Meerbeek, Arnold van den Burg and Erwin Al for their constructive feedback. E.D. holds a SB-doctoral fellowship of the Research Foundation Flanders (FWO, 1S43617N).

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.foreco.2020.118377.

References

- Aerts, R., Ewald, M., Nicolas, M., Piat, J., Skowronek, S., Lenoir, J., Hattab, T., Garzón-López, C.X., Feilhauer, H., Schmidtlein, S., Rocchini, D., Decocq, G., Somers, B., Van De Kerchove, R., Denef, K., Honnay, O., 2017. Invasion by the alien tree prunus serotina alters ecosystem functions in a temperate deciduous forest. Front. Plant Sci. 8, 1–11. https://doi.org/10.3389/fpls.2017.00179.
- Bowman, W.D., Cleveland, C.C., Halada, L., Hreško, J., Baron, J.S., 2008. Negative impact of nitrogen deposition on soil buffering capacity. Nat. Geosci. 1, 767–770. https:// doi.org/10.1038/ngeo339.
- Brandtberg, P.-O., Lundkvist, Heléne, Bengtsson, Jan, 2000. Changes in forest-floor chemistry caused by a birch admixture in Norway spruce stands. For. Ecol. Manage. 130 (1–3), 253–264. https://doi.org/10.1016/S0378-1127(99)00183-8.
- Brown, N., Vanguelova, E., Parnell, S., Broadmeadow, S., Denman, S., 2018. Predisposition of forests to biotic disturbance: Predicting the distribution of Acute Oak Decline using environmental factors. For. Ecol. Manage. 407, 145–154. https:// doi.org/10.1016/j.foreco.2017.10.054.
- Buurman, P., Pape, T., Reijneveld, J.A., de Jong, F., van Gelder, E., 2001. Laser-diffraction and pipette-method grain sizing of Dutch sediments: Correlations for fine fractions of marine, fluvial, and loess samples. Geol. en Mijnbouw/Netherlands J. Geosci. 80, 49–57. https://doi.org/10.1017/s0016774600022319.
- Carnol, M., Bazgir, M., 2013. Nutrient return to the forest floor through litter and throughfall under 7 forest species after conversion from Norway spruce. For. Ecol. Manage. 309, 66–75. https://doi.org/10.1016/j.foreco.2013.04.008.
- Chadwick, Oliver A., Chorover, Jon, 2001. The chemistry of pedogenic thresholds. Geoderma 100 (3–4), 321–353. https://doi.org/10.1016/S0016-7061(01)00027-1.
- Contreras, M.A., Affleck, D., Chung, W., 2011. Evaluating tree competition indices as predictors of basal area increment in western Montana forests. For. Ecol. Manage. 262, 1939–1949. https://doi.org/10.1016/j.foreco.2011.08.031.
- Dassonville, N., Vanderhoeven, S., Vanparys, V., Hayez, M., Gruber, W., Meerts, P., 2008. Impacts of alien invasive plants on soil nutrients are correlated with initial site conditions in NW Europe. Oecologia 157, 131–140. https://doi.org/10.1007/s00442-008-1054-6.
- de Vries, W., Dobbertin, M.H., Solberg, S., van Dobben, H.F., Schaub, M., 2014. Impacts of acid deposition, ozone exposure and weather conditions on forest ecosystems in Europe: an overview. Plant Soil 380, 1–45. https://doi.org/10.1007/s11104-014-2056-2.
- Demchik, M.C., Sharpe, W.E., 2000. The effect of soil nutrition, soil acidity and drought on northern red oak (Quercus rubra L.) growth and nutrition on Pennsylvania sites with high and low red oak mortality. For. Ecol. Manage. 136, 199–207. https://doi. org/10.1016/S0378-1127(99)00307-2.
- Denman, S., Kirk, S., Webber, J., 2010. Managing acute oak decline. Pract. Note For. Comm., pp. 6.
- Desie, E., Vancampenhout, K., Heyens, K., Hlava, J., Verheyen, K., Muys, B., 2019. Forest conversion to conifers induces a regime shift in soil process domain affecting carbon stability. Soil Biol. Biochem. 136, 107540. https://doi.org/10.1016/j.soilbio.2019. 107540.
- Desie, E., Vancampenhout, K., Nyssen, B., van den Berg, L., Weijters, M., van Duinen, G.-J., den Ouden, J., Van Meerbeek, K., Muys, B., 2020. Litter quality and the law of the most limiting: Opportunities for restoring nutrient cycles in acidified forest soils. Sci. Total Environ. 699, 134383. https://doi.org/10.1016/J.SCITOTENV.2019.134383.
- Driessen, P.M., 2001. Food and Agriculture Organization of the United Nations., International Soil Reference and Information Centre., International Institute for Aerospace Survey and Earth Sciences., Katholieke Universiteit te Leuven (1970-), Landbouwuniversiteit Wageningen. Lecture notes on the major soils of the world. Food and Agriculture Organization of the United Nations.
- Dumitru, I., Zdrilic, A., Azzopardi, A., 1999. Soil remineralisation with basaltic rock dust in Australia.
- FAO, 2006. Guidelines for soil description, fourth ed. Rome.
- Finzi, A.C., Canham, C.D., Breemen, N.V., 1998. Canopy tree-soil interactions within temperate forests: species effects on pH and cations. Ecol. Appl. 8, 447–454.
- Formánek, P., Vranová, V., 2002. A contribution to the effect of liming on forest soils: Review of literature. Artic. J. For. Sci. 48, 182–190. https://doi.org/10.17221/4692-JFS.
- Galloway, J.N., 2001. Acidification of the world: natural and anthropogenic. Water, Air Soil Pollut. 130, 17–24.
- Grasshoff, K., Johannsen, H., 1977. A new sensitive method for the determination of ammonia in sea water. Water Res. 2.

- Gruba, P., Mulder, J., 2015. Tree species affect cation exchange capacity (CEC) and cation binding properties of organic matter in acid forest soils. Sci. Total Environ. 511, 655–662. https://doi.org/10.1016/J.SCITOTENV.2015.01.013.
- Hättenschwiler, S., Tiunov, A.V., Scheu, S., 2005. Biodiversity and Litter Decomposition in Terrestrial Ecosystems. Annu. Rev. Ecol. Evol. Syst. 36, 191–218. https://doi.org/ 10.1146/annurev.ecolsys.36.112904.151932.
- Hobbie, S.E., Ogdahl, M., Reich, P.B., Oleksyn, J., Hale, C., Zytkowiak, R., Karolewski, P., 2006. Tree species effects on decomposition and forest floor dynamics in a common garden. Ecology 87, 2288–2297. https://doi.org/10.1890/0012-9658(2006) 87[2288:TSEODA]2.0.CO;2.
- Hommel, P., de Waal, R., de; Muys, B., den Ouden, J., Spek, T., 2007. Terug naar het lindewoud : strooiselkwaliteit als basis voor ecologisch bosbeheer. KNVV Uitgeverij, Zeist.
- Hüttl, R.F., Schneider, B.U., 1998. Forest ecosystem degradation and rehabilitation. Ecol. Eng. 10, 19–31. https://doi.org/10.1016/S0925-8574(97)10023-4.
- IUSS Working Group WRB, 2015. World Reference Base for Soil Resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps. Rome.
- Jonard, M., Andre, F., Ponette, Q., 2006. Modeling leaf dispersal in mixed hardwood forests using a ballistic approach. Ecology 87, 2306–2318. https://doi.org/10.1890/ 0012-9658(2006)87[2306:MLDIMH]2.0.CO;2.
- Jung, T., Blaschke, H., Obwald, W., 2000. Involvement of soilborne Phytophthora species in central European oak decline and the effect of site factors on the disease. Plant Pathol. 49, 706–718. https://doi.org/10.1046/j.1365-3059.2000.00521.x.
- Kasse, C., Vandenberghe, D., De Corte, F., 2007. e Weichselian fluvio-aeolian sands and coversands of the type locality Grubbenvorst (southern Netherlands): sedimentary environments, climate record and age. J. Quat. Sci. 22, 695–708.
- Kingston, Haswell, 1997. Microwave-Enhanced Chemistry Fundamentals, Sample Preparation, and Applications. Washington.
- Kreutzer, K., 1995. Effects of forest liming on soil processes, in: Nutrient Uptake and Cycling in Forest Ecosystems. Springer Netherlands, Dordrecht, pp. 447–470. https:// doi.org/10.1007/978-94-011-0455-5_51.
- Lavelle, P., Charpentier, F., Villenave, C., Rossi, J.-P., Derouard, L., Pashanasi, B., André, J., Ponge, J.-F., Bernier, N., Clive, A.E., 2004. Effect of earthworms on soil organic matter and nutrient dynamics at a landscape scale over decades. Earthworm Ecol. 145–160. https://doi.org/10.1201/9781420039719.pt4.
- Lavelle, P., Dugdale, R., Scholes, R., Asefaw Berhe, A., Carpenter, E., Codispoti, L., Izac, A.-M., Lemoalle, J., Luizao, F., Scholes, M., Tréguer, P., Ward, B., 2003. Ecosystems and human well-being: a report of the Millennium Ecosystem Assessment: Nutrient Cycling (Chapter 12).
- Likens, G.E., Driscoll, C.T., Buso, D.C., 1996. Long-Term Effects of Acid Rain: Response and Recovery of a Forest Ecosystem. Science (80-.). 272, 244–246. https://doi.org/ 10.1126/science.272.5259.244.
- Liu, C.L., Wang, M.K., Yang, C.C., 2001. Determination of cation exchange capacity by one-step soil leaching column method. Commun. Soil Sci. Plant Anal. 32, 15–16. https://doi.org/10.1081/CSS-120000378.
- Lorenz, K., Preston, C.M., Krumrei, S., Feger, K.-H., 2004a. Decomposition of needle/leaf litter from Scots pine, black cherry, common oak and European beech at a conurbation forest site. Eur. J. For. Res. 123, 177–188. https://doi.org/10.1007/s10342-004-0025-7.
- Mueller, K.E., Hobbie, S.E., Chorover, J., Reich, P.B., Eisenhauer, N., Castellano, M.J., Chadwick, O.A., Dobies, T., Hale, C.M., Jagodziński, A.M., Jagodziński, J., Kałucka, I., Kieliszewska-Rokicka, B., Modrzyński, J., Modrzyński, M., Ro, Zen •, A., Skorupski, M., Łukasz, •, Małgorzata, S.•, Stasińska, S., Trocha, L.K., Weiner, J., Wierzbicka, A., Oleksyn, J., Mueller, K.E., Hobbie, S.E., Chorover, J., Reich, P.B., Oleksyn, Á.J., Eisenhauer, N., Castellano, M.J., Chadwick, O.A., 2015. Effects of litter traits, soil biota, and soil chemistry on soil carbon stocks at a common garden with 14 tree species 123, 313–327. https://doi.org/10.1007/s10533-015-0083-6.
- Musil, I., Pavlicek, V., 2002. Liming of forest soils: effectiveness of particle-size fractions. J. For. Sci. 48, 121–129.
- Muys, B., Lust, N., Granval, P., 1992. Effects of grassland afforestation with different tree species on earthworm communities, litter decomposition and nutrient status. Soil Biol. Biochem. 24, 1459–1466. https://doi.org/10.1016/0038-0717(92)90133-I.
- Nickmans, H., Jonard, M., Verheyen, K., Ponette, Q., 2019. Modelling leaf dispersal and nutrient return in tree species mixtures. For. Ecol. Manage. 436, 68–78. https://doi. org/10.1016/J.FORECO.2019.01.001.
- Paquette, A., Hector, A., Castagneyrol, B., Vanhellemont, M., Koricheva, J., Scherer-Lorenzen, M., Verheyen, K., 2018. A million and more trees for science. Nat. Ecol. Evol. 2, 763–766. https://doi.org/10.1038/s41559-018-0544-0.
- Peterken, G.F., George, F., 1996. Natural woodland: ecology and conservation in northern temperate regions. Cambridge University Press.
- Ponge, J.-F., Chevalier, R., Loussot, P., 2002. Humus Index: an integrated tool for the assessment of forest floor and topsoil properties. Soil Sci. Soc. Am. J. 66, 1996. https://doi.org/10.2136/sssaj2002.1996.
- Qian, Song S., Cuffney, Thomas F., Alameddine, Ibrahim, McMahon, Gerard, Reckhow, Kenneth H., 2010. On the application of multilevel modeling in environmental and ecological studies. Ecology 91 (2), 355–361. https://doi.org/10.1890/09-1043.1.
- Reich, P.B., Oleksyn, J., Modrzynski, J., Mrozinski, P., Hobbie, S.E., Eissenstat, D.M., Chorover, J., Chadwick, O.A., Hale, C.M., Tjoelker, M.G., 2005a. Linking litter calcium, earthworms and soil properties: A common garden test with 14 tree species. Ecol. Lett. 8, 811–818. https://doi.org/10.1111/j.1461-0248.2005.00779.x.
- Schaberg, P.G., Dehayes, D.H., Hawley, G.J., 2001. Anthropogenic calcium depletion: A unique threat to forest ecosystem health? Ecosyst. Heal. 7, 214–228. https://doi.org/ 10.1046/j.1526-0992.2001.01046.x.
- Schmitz, A., Sanders, T.G.M., Bolte, A., Bussotti, F., Dirnböck, T., Johnson, J., Peñuelas, J., Pollastrini, M., Prescher, A.-K., Sardans, J., Verstraeten, A., de Vries, W., 2019.

Responses of forest ecosystems in Europe to decreasing nitrogen deposition. Environ. Pollut. 244, 980–994. https://doi.org/10.1016/J.ENVPOL.2018.09.101.

- Schulte, E.E., Hopkins, B.G., 1996. Estimation of organic matter by weight loss-on-ignition. In: Magdoff, F.R., et al. (Ed.), Soil Organic Matter: Analysis and Interpretation. SSSA Spec. Pub. No. 46, Madison, pp. 21–31.
- Starfinger, U., Kowarik, I., Rode, M., Schepker, H., 2003. From desirable ornamental plant to pest to accepted addition to the flora?-the perception of an alien tree species through the centuries. Biol. Invas.
- Ste-Marie, C., Paré, D., 1999. Soil, pH and N availability effects on net nitrification in the forest floors of a range of boreal forest stands. Soil Biol. Biochem. 31, 1579–1589. https://doi.org/10.1016/S0038-0717(99)00086-3.
- Team, R.C., 2019. R: A language and environment for statistical computing. R Found. Stat. Comput.
- Technicon, 1969. Industrial method 33–69W, nitrate + nitrite in water. In: Technicon Autoanalyser Methodology. Technicon Corporation, Karrytown.
- Ulrich, B., Sumner, M.E., 1991. Soil Acidity. Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-74442-6.
- Van Nevel, L., Mertens, J., De Schrijver, A., De Neve, S., Verheyen, K., 2014. Can shrub species with higher litter quality mitigate soil acidification in pine and oak forests on poor sandy soils? For. Ecol. Manage. 330, 38–45. https://doi.org/10.1016/j.foreco. 2014.07.002.
- Van Ranst, E., Sys, C., 2000. Eenduidige legende voor de digitale bodemkaart van Vlaanderen (schaal 1:20 000). Gent.
- Vanderhoeven, S., Dassonville, N., Meerts, P., 2005. Increased Topsoil Mineral Nutrient Concentrations Under exotic invasive plants in Belgium. Plant Soil 275, 169–179. https://doi.org/10.1007/s11104-005-1257-0.
- Vannoppen, A., Treydte, K., Boeckx, P., Kint, V., Ponette, Q., Verheyen, K., Muys, B.,

- 2020. Tree species diversity improves beech growth and alters its physiological response to drought. Trees 1–15. https://doi.org/10.1007/s00468-020-01981-0.
- Verstraeten, G., Vancampenhout, K., Desie, E., De Schrijver, A., Hlava, J., Schelfhout, S., Verheyen, K., Muys, B., 2018. Tree species effects are amplified by clay content in acidic soils. Soil Biol. Biochem. 121, 43–49. https://doi.org/10.1016/j.soilbio.2018. 02.021.
- Vitousek, P.M., Chadwick, O.A., 2013. Pedogenic Thresholds and Soil Process Domains in Basalt-Derived Soils. Ecosystems 16, 1379–1395. https://doi.org/10.1007/s10021-013-9690-z.
- von Wendorff, G., 1952. Die Prunus serotina in Mitteleuropa. Eine waldbauliche Monographie. Universität Hamburg.
- Waldner, P., Thimonier, A., Graf Pannatier, E., Etzold, S., Schmitt, M., Marchetto, A., Rautio, P., Derome, K., Nieminen, T.M., Nevalainen, S., Lindroos, A.-J., Merilä, P., Kindermann, G., Neumann, M., Cools, N., de Vos, B., Roskams, P., Verstraeten, A., Hansen, K., Pihl Karlsson, G., Dietrich, H.-P., Raspe, S., Fischer, R., Lorenz, M., Iost, S., Granke, O., Sanders, T.G.M., Michel, A., Nagel, H.-D., Scheuschner, T., Simončič, P., von Wilpert, K., Meesenburg, H., Fleck, S., Benham, S., Vanguelova, E., Clarke, N., Ingerslev, M., Vesterdal, L., Gundersen, P., Stupak, I., Jonard, M., Potočić, N., Minaya, M., 2015. Exceedance of critical loads and of critical limits impacts tree nutrition across Europe. Ann. For. Sci. 72, 929–939. https://doi.org/10.1007/ s13595-015-0489-2.
- Zanella, A., Jabiol, B., Ponge, J.F., Sartori, G., De Waal, R., Van Delft, B., Graefe, U., Cools, N., Katzensteiner, K., Hager, H., Englisch, M., Brethes, A., Broll, G., Gobat, J. M., Brun, J.J., Milbert, G., Kolb, E., Wolf, U., Frizzera, L., Galvan, P., Koll, R., Baritz, R., Kemmers, R., Vacca, A., Serra, G., Banas, D., Garlato, A., Chersich, S., Klimo, E., Langohr, R., 2014. European Humus Forms Reference Base.